

Splitting Enables Overcoming the Curse of Dimensionality

Jerome Darbon* Stanley Osher†

In this short note we briefly outline a new and remarkably fast algorithm for solving a large class of high dimensional Hamilton-Jacobi (H-J) initial value problems arising in optimal control and elsewhere [1]. This is done without the use of grids or numerical approximations. Moreover, by using the level set method [8] we can rapidly compute projections of a point in \mathbb{R}^n , n large to a fairly arbitrary compact set [2]. The method seems to generalize widely beyond what we will present here to some nonconvex Hamiltonians, state dependent Hamiltonians, differential games and perhaps new linear programming algorithms.

We begin with the HJ initial value problem

$$\begin{cases} \frac{\partial \varphi}{\partial t}(x, t) + H(\nabla_x \varphi(x, t)) = 0 & \text{in } \mathbb{R}^n \times (0, +\infty) \\ \varphi(x, 0) = J(x) & \forall x \in \mathbb{R}^n. \end{cases} \quad (1)$$

We assume $J : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex and one coercive, i.e., $\lim_{\|x\|_2 \rightarrow +\infty} \frac{J(x)}{\|x\|_2} = +\infty$, $H : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex and positively one homogeneous (we sometimes relax all these assumptions).

A good example of this is

$$H(v) = \|v\|_2.$$

Here $\|v\|_p = (\sum_{i=1}^n |v_i|^p)^{\frac{1}{p}}$ for $p \geq 1$ and $\langle x, v \rangle = \sum_{i=1}^n x_i v_i$.

If we take for J a convex Lipschitz function having the property that, for Ω a convex compact set of \mathbb{R}^n

$$\begin{cases} J(x) < 0 & \text{for any } x \in \text{int } \Omega \\ J(x) = 0 & \text{for any } x \in (\Omega \setminus \text{int } \Omega) \\ J(x) > 0 & \text{for any } x \in (\mathbb{R}^n \setminus \Omega). \end{cases}$$

*CNRS/CMLA-Ecole Normale Supérieure de Cachan

†Mathematics, UCLA, Los Angeles, CA 90095-1555, Research supported by ONR grants N000151510683, N000141210838 and DOE grant DE-SC00183838

We call this level set initial data. Then the set of points for which $\varphi(x, t) = 0$, $t > 0$ are exactly those at a distance t , from the boundary of Ω . In fact given $\bar{x} \notin \Omega$, then the closest point x_{opt} from \bar{x} to $\partial\Omega$ is exactly

$$x_{\text{opt}} = \bar{x} - t \frac{\nabla\varphi(\bar{x}, t)}{\|\nabla\varphi(\bar{x}, t)\|_2}. \quad (2)$$

To solve (1) we use the Hopf formula [5]

$$\varphi(x, t) = (J^* + tH)^*(x) = - \min_{v \in \mathbb{R}^n} \{J^*(v) + tH(v) - \langle x, v \rangle\},$$

where the Fenchel-Legendre transform $f^* : \mathbb{R}^n \rightarrow R \cup (+\infty)$ of the convex function f is defined by

$$f^*(v) = \sup_{x \in \mathbb{R}^n} \{\langle v, x \rangle - f(x)\}.$$

Moreover, for free we get that the minimizer satisfies

$$\arg \min_{v \in \mathbb{R}^n} \{J^*(v) + tH(v) - \langle x, v \rangle\} = \nabla_x \varphi(x, t). \quad (3)$$

whenever $\varphi(\cdot, t)$ is differentiable at x . Let us note here that our algorithm computes $\varphi(x, t)$ but also $\nabla_x \varphi(x, t)$.

Also, we can use the Hopf-Lax formula [5, 6] to solve (1).

$$\varphi(x, t) = \min_{z \in \mathbb{R}^n} \left\{ J(z) + tH^* \left(\frac{x - z}{t} \right) \right\} \quad (4)$$

for convex H .

From (4) it is easy to show that if we have k different initial value problems $i = 1, \dots, k$

$$\begin{cases} \frac{\partial \varphi_i}{\partial t}(x, t) + H(\nabla_x \varphi_i(x, t)) = 0, & \text{in } \mathbb{R}^n \times (0, +\infty) \\ \varphi_i(x, 0) = J_i(x) & \forall x \in \mathbb{R}^n \end{cases}$$

with the usual hypotheses, then (4) implies, for any $x \in \mathbb{R}^n$, $t > 0$

$$\varphi_i(x, t) = \min_{z \in \mathbb{R}^n} \left\{ J_i(z) + tH^* \left(\frac{x - z}{t} \right) \right\}.$$

So

$$\min_{i=1, k} \varphi_i(x, t) = \min_{z \in \mathbb{R}^n} \left\{ \min_{i=1, \dots, k} \left\{ J_i(z) + tH^* \left(\frac{x - z}{t} \right) \right\} \right\}$$

solves the initial value problem

$$\begin{cases} \frac{\partial \varphi}{\partial t}(x, t) + H(\nabla_x \varphi(x, t)) = 0, & \text{in } \mathbb{R}^n \times (0, +\infty) \\ \varphi(x, 0) = \min_{i=1, \dots, k} J_i(x) & \forall x \in \mathbb{R}^n. \end{cases} \quad (5)$$

This means that if $\Omega = \cup_{i=1, \dots, k} \Omega_i$, each Ω_i is compact and convex and may overlap, then we can easily compute the set of all points at distance t from Ω which is exactly the solution to (5) where each J_i is a level set function for Ω_i . Moreover, at every point \bar{x} outside of $\bar{\Omega}$ for which there is one i such that $\varphi_i(\bar{x}, t) < \varphi_{i'}(\bar{x}, t)$ for any $i \neq i'$, then the closest point x_{opt} to \bar{x} and Ω is again

$$x_{\text{opt}} = \bar{x} - t \frac{\nabla_x \varphi_i(\bar{x}, t)}{|\nabla_x \varphi_i(\bar{x}, t)|}.$$

If there are several i for which $\varphi_i(\bar{x}, t)$ is the minimum among all k of them, then $\nabla_x \varphi$ will be “multivalued”, i.e. it will have jumps, but any of the x_{opt} defined above will be a closest point on Ω to \bar{x} .

We solve the optimization problem (3) by using the split Bregman algorithm [4, 3, 9] as follows

$$v^{k+1} = \arg \min_{v \in \mathbb{R}^n} \{J^*(v) - \langle x, v \rangle + \frac{\lambda}{2} \|d^k - v - b^k\|_2^2\}, \quad (6)$$

$$d^{k+1} = \arg \min_{d \in \mathbb{R}^n} \left\{ tH(d) + \frac{\lambda}{2} \|d - v^{k+1} - b^k\|_2^2 \right\} \quad (7)$$

$$b^{k+1} = b^k + v^{k+1} - d^{k+1}. \quad (8)$$

Here the sequences $(v^k)_{k \in \mathbb{N}}, (d^k)_{k \in \mathbb{N}}$ both converge to $\nabla_x \varphi(x, t)$. Let us emphasize again that our numerical algorithm not only computes the solution $\varphi(x, t)$ but also computes $\nabla_x \varphi(x, t)$ when $\varphi(\cdot, t)$ is differentiable.

Both (6) and (7), up to change of variables, can be reformulated as finding the unique minimizer of

$$\arg \min_w \left\{ \alpha f(w) + \frac{1}{2} \|w - z\|_2^2 \right\}$$

which is the proximal map of f . Equation (6) can be solved if either J^* or J have easily computable proximal maps, which often occurs, especially if one of them is smooth.

Equation (7) can be easily solved if $H(d) = \|d\|_2$ via the shrink₂ operator defined by

$$\text{shrink}_2(z, \alpha) = \begin{cases} \frac{z}{\|z\|_2} \max(\|z\|_2 - \alpha, 0) & \text{if } z \neq 0 \\ 0 & \text{if } z = 0 \end{cases}$$

and we have

$$\arg \min_w \left\{ \alpha \|w\|_2 + \frac{1}{2} \|w - z\|_2^2 \right\} = \text{shrink}_2(z, \alpha)$$

If $H(d) = \|d\|_1$ we use shrink_1 operator defined as follows for any $i = 1, \dots, n$

$$(\text{shrink}_1(z, \alpha))_i = \begin{cases} z_i - \alpha & \text{if } z_i > \alpha \\ 0 & \text{if } |z_i| \leq \alpha \\ z_i + \alpha & \text{if } z_i < -\alpha \end{cases}$$

and we have

$$\arg \min_w \left\{ \alpha \|w\|_1 + \frac{1}{2} \|w - z\|_2^2 \right\} = \text{shrink}_1(z, \alpha).$$

To solve (7) for more general $H(d)$ convex one homogeneous or to find the proximal map for f of that type we use the fact that H^* is the characteristic function of a closed convex set $C \subset \mathbb{R}^n$

$$H^* = I_C.$$

By using the Moreau identity [7] we realize that the proximal map of H can be obtained by projecting onto C . To do this projection, we merely solve the eikonal equation with level set initial data for C via split Bregman as above in (6), (7), (8) with $H(d) = \|d\|_2$. This is easy using the shrink_2 operator. We then use (2) to obtain the projection and repeat the entire iteration.

Numerical experiments on an Intel Laptop Core i5-5300U running at 2.3 GHz are now presented. We consider diagonal matrices D defined by $D_{ii} = 1 + \frac{1+i}{n}$ for $i = 1, \dots, n$. We also consider matrices A defined by $A_{ii} = 2$ for $i = 1, \dots, n$ and $A_{ij} = 1$ for $i, j = 1, \dots, n$. Table 1 presents the average time (in seconds) to evaluate the solution over 1,000,000 samples (x, t) uniformly drawn in $[-10, 10]^n \times [0, 10]$. The convergence is remarkably rapid: 10^{-6} to 10^{-8} seconds on a standard laptop, per function evaluation. Figure 1 depicts 2-dimensional slices at different times for the (H-J) equation with a weighted ℓ_1 Hamiltonian $H = \|D \cdot\|_1$, initial data $J = \frac{1}{2} \|\cdot\|_2^2$ and $n = 8$.

References

- [1] J. Darbon and S. Osher, *Algorithms for Overcoming the Curse of Dimensionality for Certain Hamilton-Jacobi Equations Arising in Control Theory and Elsewhere* UCLA CAM report 15-50.

n	$\ y\ _1$	$\ y\ _2$	$\ y\ _\infty$	$\ y\ _D$	$\ y\ _A$
4	6.36e-08	1.20e-07	2.69e-07	7.00e-07	8.83e-07
8	6.98e-08	1.28e-07	4.89e-07	1.07e-06	1.57e-06
12	8.72e-08	1.56e-07	7.09e-07	1.59e-06	2.23e-06
16	9.24e-08	1.50e-07	9.92e-07	2.04e-06	2.95e-06

Table 1: Time results in seconds for the average time per call for evaluating the solution of the HJ-PDE with the initial data $J = \frac{1}{2}\|\cdot\|_2^2$, several Hamiltonians and various dimensions n .

- [2] J. Darbon and S. Osher, *Fast projections onto compact sets in high dimensions using the level set method, Hopf formulas and optimization*. To appear.
- [3] R. Glowinski and A. Marrocco, *Sur l'approximation par éléments finis d'ordre, et la résolution par pénalisation-dualité, d'une classe de problèmes de Dirichlet non-linéaires*, C.R. hebdomadaire Séances Acad. Sci., Paris 278, série A, (1974), pp. 1649-1652.
- [4] T. Goldstein and S. Osher, *The Split Bregman Method for L_1 Regularized Problems*, SIAM J. on Imaging Science, v.2, (2), 2009, pp. 323-343.
- [5] E. Hopf, *Generalized Solutions of Nonlinear Equations of the First Order*, J. Math. Mech., v.14, (1965), pp. 951-973.
- [6] P. D. Lax, *Hyperbolic Systems of Conservation Laws and the Mathematical Theory of Conservation Laws*, SIAM, 1973
- [7] J.-J. Moreau. *Proximité et dualité dans un espace hilbertien*. Bulletin Soc. Math. France, 93, pp. 273–299, 1965.
- [8] S. Osher and J.A. Sethian, *Fronts Propagating with Curvature Dependent Speed: Algorithms Based on Hamilton-Jacobi Formulations*, J. Comput. Phys., v.79, (1), (1988), pp. 12-49.
- [9] S. Osher and W. Yin, *Error Forgetting of Bregman Iteration*, J. Sci. Comput., v.54, (2), (2013), pp. 684-695.




Figure 1: Evaluation of the solution $\phi((x_1, x_2, 0, 0, 0, 0, 0)^\dagger, t)$ of the HJ-PDE with initial data $J = \frac{1}{2} \|\cdot\|_2^2$ and Hamiltonian $H = \|D \cdot\|_1$ for $(x_1, x_2) \in [-20, 20]^2$ for different times t . Plots for $t = 0, 3, 5, 8$ and respectively depicted in (a), (b), (c) and (d). The level lines multiple of 10 are superimposed on the plots.